




# **Environmental Impact Statement (EIS) for Permitted Use of Triclopyr**

---

**Draft**

**February 2004  
Publication Number 04-10-XXX**

 *Printed on Recycled Paper*



# **Environmental Impact Statement (EIS) for Permitted Use of Triclopyr**

---

**Draft**

Washington State Department of Ecology  
Water Quality Program

February 2004  
Publication Number 04-10-XXX



*Printed on Recycled Paper*


For additional copies of this document contact:

Department of Ecology  
Publications Distribution Center  
P.O. Box 47600  
Olympia, WA 98504-7600

Telephone: (360) 407-7472

**Headquarters (Lacey) 360-407-6000**

**If you are speech or hearing impaired, call 711 or 1-800-833-6388 for TTY**



*If you need this publication in an alternate format, please contact us at 360-407-6404).  
The TTY number number (for the speech and hearing impaired) is 711 or 1-800-833-6388.*

# Table of Contents

Table of Contents .....	i
Draft Environmental Impact Statement (EIS) for Permitted Use of Triclopyr .....	1
1. Registration Status .....	1
2. Description .....	2
3. Environmental and Human Health Impacts .....	5
Air .....	5
Earth .....	7
Water .....	8
Estuarine (Intertidal) Environments .....	9
Plants .....	10
Animals .....	11
Water, Land and Shoreline Use .....	16
4. Mitigation Summary for Triclopyr TEA .....	19
References .....	21



# Draft Environmental Impact Statement (EIS) for Permitted Use of Triclopyr Under Ecology's Water Quality Program

This draft EIS is a supplement to the Final Supplemental Environmental Impact Statement for Freshwater Aquatic Plant Management (Ecology, 2001). The purpose of this Supplemental Environmental Impact Statement is to review what is known about the potential environmental effects of aquatic uses of triclopyr. This includes a summary of the registration status, environmental effects, potential human health impacts, and recommended mitigation to minimize the effects of triclopyr application. This information is designed to follow closely the State Environmental Protection Act (SEPA) Checklist. Except where otherwise noted, references to the 2001 triclopyr risk assessment prepared for Ecology by Compliance Services International (CSI) Volume 5 Triclopyr, Sections 1 – 5. Mitigations for the use of triclopyr in Washington waters that go beyond label conditions are bulleted in each section and summarized at the end of this document.

## 1. Registration Status

Triclopyr TEA (triethylamine) was first registered by Dow AgroSciences on May 8, 1979 as an herbicide on non-crop areas and in forestry use for the control of broadleaf weeds and woody plants. In 1984, it was registered for use on turf sites. In 1995, triclopyr TEA was registered for use on rice to control many hard to control broadleaf weed species. EPA's Re-registration Eligibility Decision (RED) process for triclopyr acid, triclopyr TEA and triclopyr BEE (butoxyethyl ester) was completed on September 30, 1997. The on-line RED can be viewed at <http://www.epa.gov/oppsrrd1/REDs/2710red.pdf>.

Garlon® 3A (EPA Reg. No. 62719-37) from Dow AgroScience is currently registered in the state of Washington for the control of aquatic weeds in public water ways and annual and perennial broadleaf weeds and woody brush in wetlands.

November 2002, SePRO Corporation received Federal EPA registration for Triclopyr TEA salt under the trade name (EPA Reg. No. 62719-37-67690). The Renovate® label specifies selective control of nuisance and exotic plants such as Eurasian watermilfoil (*Myriophyllum spicatum*), purple loosestrife (*Lythrum salicaria*), alligatorweed (*Alternanthera philoxeroides*), and water hyacinth (*Eichhornia crassipes*).

DowElanco currently manufactures and distributes Garlon® 3A and SEPRO Corporation will market and distribute Renovate® under a separate label. The products will be the same since DowElanco will manufacture both products.

## 2. Description

### *What triclopyr is used for*

Triclopyr, ((3,5,6-trichloro-2-pyridinyl) oxyacetic acid) is an aquatic herbicide that utilizes a systemic mode of action used to control submerged, floating and emergent aquatic plants in both static and flowing water. It is also registered for a number of terrestrial uses including broadleaf weed control, and is used in rice, pasture and rangeland, rights-of-way, forestry, turf, and home lawns and gardens.

### *Other ingredients in the triclopyr formulation*

Triclopyr is formulated as a solution in water. Intentionally added inert or “other” ingredients in triclopyr formulations include water and triethanol amine (TEA). The water serves as the primary diluent/solvent in the liquid product while the triethanol amine is used to form the salt of the technical grade active ingredient. There are no known impurities identified by the manufacturers or the US EPA that are known to be of toxicological or environmental concern.

### *How triclopyr works*

Triclopyr is a growth hormone of the *auxin* type. An auxin-type herbicide interferes with growth after the plant emerges. It contacts leaves, where sugar is produced, and moves to roots, tips, and parts of the plant that store energy, thereby interrupting growth. Since the movement of sugars from the leaves to other parts of the plant is essential for growth, this type of herbicide has the potential to kill simple perennial and creeping perennial weeds with only one or two foliar applications. Bending and twisting of leaves and stems is evident almost immediately after application. Delayed symptom development includes root formation on dicot stems; misshapen leaves, stems and flowers; and abnormal roots (EPA, 1998) (Purdue, 2000).

### *What target aquatic plants triclopyr affects*

Triclopyr has been claimed to be effective for a variety of fully or partially aquatic plants including American lotus (*Nelumbo lutea*), Eurasian watermilfoil (*Myriophyllum spicatum*), parrotfeather (*Myriophyllum aquaticum*), pennywort (*Hydrocotyle* spp.), waterhyacinth (*Eichhornia crassipes*), water lilies (*Nuphar* spp. and *Nymphaea odorata*) and waterprimrose (*Ludwigia uruguayensis*), alligatorweed (*Alternanthera philoxeroides*) and purple loosestrife (*Lythrum salicaria*).

### *What aquatic plants triclopyr does not affect*

Triclopyr is not typically used for algae control and most species of algae are not affected strongly by triclopyr (Section 4, Tables 2, 11 and 16). Many species of native plants are



not affected by triclopyr or are not affected except transitorily. Some of these may include pondweed species and coontail, rushes and cattails (Petty et al, 1998). However, at higher use rates (2.5 ppm a.e.), the more susceptible native species such as coontail, Southern naiad, and American waterweed may be reduced in numbers in some treatment situations.

#### *Physical and chemical characteristics of triclopyr*

Triclopyr dissolves readily in water but is not volatile. It has a low vapor pressure ( $1.26 \times 10^{-6}$  mm/Hg at 25° C for the active ingredient triclopyr), and a low distribution coefficient (0.165 to 0.925 mL/g). *Hydrolysis* refers to the chemical interaction of the chemical with water as a mechanism of chemical breakdown. Triclopyr acid, which forms immediately when Triclopyr TEA dissolves in water, is not hydrolyzed.

#### *Degradation mechanisms and products*

The main degradation products of triclopyr in the environment occur because of photolysis or microbial degradation.

- *Photolysis*

Photolysis, or chemical breakdown due to sunlight, can contribute substantially to the degradation of triclopyr acid and triclopyr TEA. These triclopyr products are degraded rapidly under natural sunlight (0.6 to 6.6 days) with both the dominant degradate and degradation rate varying somewhat with the product tested (EPA RED, 1998).

In sterile buffered water, photolysis of triclopyr acid (dissociated triclopyr TEA) produced MDPA (5-chloro-3, 6-dihydroxy-2-pyridinyloxyacetic acid) with small amounts of oxamic acid and carbon dioxide [EPA RED (1995) p. 53 and Woodburn et al (1993)]. In natural river water, photolysis produced mainly oxamic acid.

- *Microbial Degradation*

In aerobic soils, triclopyr degrades to carbon dioxide. Intermediate degradation products include TCP (*3,5,6-trichloro-2-pyridinol*) and TMP (*3,5,6 trichloro-2- methoxypyridine*).

#### *Persistence*

The persistence of triclopyr and its degradates varies widely depending on the conditions of the system being tested. For the most part, triclopyr is dissipated rapidly from the water column and is not adsorbed on sediment for very long periods of time.

In soils, factors that effect persistence of triclopyr include temperature, pH, higher organic matter content, higher microbial numbers, and the presence of triclopyr due to

previous applications. Half-life persistence can range from less than one day to nearly a year (Section 3, Table 3.4, p. 31). A Dow AgroSciences product (trade name Confront, containing 33% Triclopyr TEA) was not broken down by composting operations produced at a publicly owned solid waste facility in Spokane County, Washington, where it tainted at least 47,000 cubic yards of compost. The presence of the herbicide Confront rendered the compost damaging to crops and resulted several claims to the waste facility. It is unknown whether triclopyr or the herbicide's other active ingredient, clopyralid triethylamine salt (12.1%) was at fault for contaminating the compost (See: County wants Dow Chemical to stop distributing herbicide, The Spokesman Review, Spokane, WA, 4/25/01).

#### *Persistence in water*

The environmental persistence of triclopyr products in the field can be quite variable; the dissipation half-life in water varies from less than 1 day to approximately 7.5 days. However, according to most authors, the most typical half-life would be between 3.5 and 7.5 days (CSI, Volume 5, Section 3, Table 3.5, p. 55). Dissipation of triclopyr is primarily due to photolysis, degradation by microbes, and mixing of triclopyr treated water with water that has not been treated.

Dissipation is related to lake size, wind, and the amount of water exchange that occurs. The larger the lake, the more wind blowing across the lake surface, the more water exchange through inlet and outlet streams or rivers, the more likely it is that triclopyr residues will be rapidly dispersed and diluted to below detection limits. In small lakes, detectable concentrations of triclopyr may be carried a significant distance down an outlet stream if the flow is sufficient and degradation is slow.

The concentration of the toxic metabolite (TCP) has generally been low in lake and pond water with concentrations of TCP not higher than ~0.1 ppm in Lake Minnetonka, Lake Seminole, and various ponds on the day of application. It generally dissipates to concentrations below the detection limit at three days after treatment.

#### *Persistence in water - anaerobic conditions*

Anaerobic environments lack oxygen. According to studies, triclopyr in anaerobic aquatic sediment may have a very long half-life. For example, it can take a year for 20% of the amount of triclopyr to degrade in these environments. The calculated rate for 50% to degrade is about 3 and 1/2 years.

#### *Persistence on sediments*

In the Lake Minnetonka study, the half-lives of triclopyr in the sediment ranged from around five or six days at Lake Minnetonka, and the sediment half lives of TCP were approximately eleven days.

#### *Persistence in aerobic soil*

Laboratory studies indicate that triclopyr acid and the dissociated triethylamine (TEA) are readily degraded in aerobic soil. The half-lives of triclopyr acid and triclopyr TEA can vary from just a few days in laboratory aerobic soil metabolism experiments to approximately two weeks on Northern Ontario forest soils to nearly three months in pastureland in Oregon (DT50 = 75 to 81 days) (SEIS, Vol. 5, Sect. 3, Table 3.4).

When the break-down products TCP and TMP were measured in the field, it is clear that TMP is not persistent and never exceeds 0.06 ppm in any soil profile on bare ground and pastureland soil. Concentrations of TCP have been observed at 0.1 to ~0.2 ppm or higher for three-quarters of a year or longer after application of triclopyr to pasturelands.

#### *Persistence in rice paddy soils at various depths*

In the laboratory, rice paddy soils yielded half-lives that varied considerably. Rates of degradation on surface soils (DT50 = 9 to 307 days) was much greater than on soils taken from depths of about one foot (DT50 = 35 to 314 days). This phenomenon was attributed to the fact that surface soils when compared to subsurface soils had higher pH, higher organic matter content, higher microbial numbers, and the presence of acclimated versus non acclimated decomposers (microbes).

### **3. Environmental and Human Health Impacts**

#### **Air**

Triclopyr products have very little tendency to affect air quality or cause crop damage because of low vapor pressure. (The vapor pressure of commercial products of triclopyr is  $1.26 \times 10^{-6}$  mm/Hg at 25° C for the active ingredient triclopyr).

Typically, the mode of application is **subsurface injection** for liquid formulations, making drift outside the treatment area unlikely.

For those cases where a **boom sprayer** applies a liquid formulation, as much as one percent of the application may drift out of the treatment area. It has been estimated for general herbicides that this amount of drift could have an impact if 120 swaths were applied and one percent of the applied pesticide drifted out of the treatment area on each pass. In this case, dosage levels higher than that intended for the target could accumulate downwind of the treatment area. This could cause an effect on non-target plants that may damage habitat and decrease the amount of forage available for waterfowl and fish in non-target areas (Forsythe et al, 1997).

For **aerial application**, as much as 17 percent of the treatment would not strike the target area. In this case, drift out of the treatment area could impinge on non-target organisms at a great distance from the site of application. Depending on how much triclopyr was deposited per unit area outside the site, there could be a significant impact on non-target

wild plants or crops. In addition to effects on plants, non-target sensitive terrestrial wildlife may be adversely impacted.

**Odor** is unlikely to be noticed except for short periods of time following application. Since there would rarely be more than one or two applications of triclopyr per water body per year in the state of Washington, any adverse impact on quality of life due to problems with odor from triclopyr applications should be weighed carefully with the impact due to the effects of poor navigability, and effects on the recreational use of the water body.

**Direct effects from breathing** the vapor are unlikely for the general public since the acute LC<sub>50</sub> for triclopyr TEA is greater than 2.6 mg/L (EPA RED, 1998). However, EPA has determined that there are potential exposures to persons involved in mixing, loading or entering treated sites after application is complete.

**Release of Toxic Materials – Inhalation** The Washington State Department of Health conducts a Pesticide Surveillance Program and has documentation of seven human exposure cases, possibly related to terrestrial use, involving skin and upper respiratory tract irritation following direct exposure to triclopyr spray mist. No signs and symptoms of systemic poisoning were reported, however some of the individuals experienced temporary irritation of the skin and upper respiratory tract. It should be noted that application of the triclopyr product Renovate® Aquatic Herbicide is directly injected under the water and not applied by aerial or spray application (WDOH, 1999).

For **aerial application**, as much as 17 percent of the treatment would not strike the target area. In this case, drift out of the treatment area could impinge on non-target organisms at a great distance from the site of application. Depending on how much triclopyr was deposited per unit area outside the site, there could be a significant impact on non-target wild plants or crops. In addition to effects on plants, non-target sensitive terrestrial wildlife may be adversely impacted.

### ***Permit Mitigations***

#### *Drift*

Follow label instructions.

#### *Aerial applications*

- Aerial applications are not permitted in Washington State waters.

#### *Odor*

No odor problems were cited in the literature.

#### *Inhalation*

Follow label instructions.

## Earth

### *Soils*

The presence of triclopyr in soil is not anticipated from aquatic treatment unless flooding occurs or the water is used for irrigation.

### *Flooding*

If a flooding incident occurs within 120 days of application, there is a potential for triclopyr to damage upland sensitive species, particularly grapes, tobacco, vegetable crops and flowers. However, the expected half-life on soils is fairly low (8 to 18 days in the laboratory and two weeks in the field). Therefore, any adverse impact due to a flooding incident is likely to be quite limited.

### *Sediment*

In typical situations where water is fairly shallow (0.3 to less than two meters), triclopyr in sediment has observed half-lives that range from less than one day at Lake Seminole to 5.8 days in Lake Minnetonka.

Due to the low distribution coefficient for triclopyr (0.165 to 0.925 mL/g), it does not bind tightly to sediment and therefore concentrations in sediment should remain low. This assumption is confirmed by results from field studies. For example, at Lake Minnetonka, concentrations of triclopyr in sediment were never higher than 0.334 ppm a.e. and dissipation to concentrations of  $\leq 0.15$  ppm was seen within 14 days after application. At Lake Seminole, triclopyr was not seen at significant concentrations ( $< 0.1$  ppm a.e.) except for the day of application where concentrations as high as 0.64 ppm a.e. were detected. Even in the pond studies, the concentration of triclopyr in sediment was very low and did not exceed 0.86 ppm a.e. during the first few days and dissipated to below the limit of quantification within four weeks.

These low levels of triclopyr in sediment indicate that the sediment quality should remain high in treated water bodies and that such sediments should pose little or no threat to benthic in-fauna.

### *Metabolites*

The toxic metabolite TCP is found at even lower concentrations than triclopyr. Both laboratory and field studies indicate the concentrations of TCP in the sediment are very low and generally do not exceed 0.16 ppm and are typically less than 0.05 ppm in lakes and ponds. TCP generally dissipates to below the level of quantification within a few days of application to lakes but may take up to six weeks to dissipate entirely from ponds.

Another metabolite (TMP) is rarely detected in lake or pond sediment and is normally detected in the water column at concentrations that do not exceed 0.01 ppm a.e. TMP is

generally considered to have no toxicological significance. Data supporting these conclusions can be found in Getsinger et al, (2000) Petty et al (1998) and Green et al (1989) (Table 5).

#### *Anaerobic conditions*

Triclopyr from the application of Garlon® 3A or Renovate® may have long half-lives in deep sediments under anaerobic conditions. In anaerobic conditions, triclopyr degrades to TCP with a half-life of about 3.5 years (Ladowski and Didlack (1984 in Petty et al, 1998).

### ***Permit Mitigations***

#### *Irrigation*

Follow label instructions.

#### *Sediments*

- Due to the possibility of anaerobic conditions in sediments, sediment monitoring is required prior to any third application of triclopyr on a site within a three-year period. Evidence of persistence of triclopyr or TCP in sediments is basis for denial of the third application.

### **Water**

#### *Surface Water*

The concentrations of triclopyr in lakes that have been spot treated generally fall below the temporary drinking water residue tolerance (0.5 ppm a.e.) within one day but in rare instances can take as long as eight days. However, the concentration of triclopyr in ponds can take three to four weeks to dissipate to concentrations below 0.5 ppm a.e. (CSI, Volume 5, Sect. 3, Table 3.5, pp. 55-67).

The concentration of the toxic metabolite (TCP) has generally been low in lake and pond water with concentrations of TCP not higher than ~0.1 ppm in Lake Minnetonka, Lake Seminole, and various ponds on the day of application and generally dissipating to concentrations below the detection limit at three days after treatment.

Fish and other aquatic organisms need oxygen to survive and treatment of dense weed areas may result in dissolved oxygen decreases due to the decomposition of dead weeds. Therefore, application of triclopyr TEA products must be limited to a portion of the water body at any one time. Typically, the entire water body is typically not treated. Only about 20% of a water body is typically treated based on areas designated for priority control.

## *Wetlands*

Because of the manner in which triclopyr products are applied, significant impact to other wetland environments is unlikely. There may be some tendency for drift into other wetland environments or a flow of water into estuarine, palustrine, riparian, lentic or lotic environments. However, it is not anticipated that the impact would be measurable due to dilution effects, as treated ponds, lakes, and canals normally flow into streams and rivers and ultimately into estuaries.

The total application of these products should not exceed 2.5 ppm a.e. for the treatment area per annual growing season. The total application of these products to control floating and emerged weeds should not exceed two gallons formulation/acre per annual growing season.

## *Estuarine (Intertidal) Environments*

Water from a stream or river containing triclopyr may flow into an estuary. However, dilution effects from the water already present in the estuary and diurnal tides should dilute triclopyr to levels where it is not significant in the water column.

## *Palustrine (Marshy) Environments*

Most immersed plants are not likely to be adversely impacted at the concentrations of triclopyr used to control fully aquatic weeds. However, floating (*Eicchornia crassipes*) and rooted submersed plants (*Myriophyllum* spp. and *Hydrocotyle* spp.), that are typically found in a palustrine environment may be affected by water that enters these areas from lakes and ponds.

It is unclear exactly how high the triclopyr concentrations must be to damage native plant species. Initial triclopyr concentrations of 2.5 ppm a.e. that remained at levels of 1.0 ppm a.e. or higher for 7 to 14 days have been known to adversely impact coontail (*Ceratophyllum* spp.), Eurasian watermilfoil (*Myriophyllum spicatum*), southern naiads (*Naja guadalupensis*), and American waterweed (*Elodea canadensis*) in water impounds (ponds) located at Elk Grove, California, Columbia, Missouri, or Lewisville, Texas (Petty et al, 1998). If these rooted macrophytes were destroyed due to herbicide applications, there would be less tendency for the marsh to flood, resulting in loss of habitat for fish, amphibians, wild birds and mammals.

## ***Permit Mitigations***

### *Surface water*

Follow label directions.

## *Wetlands*

- The total application should not exceed 2.5 ppm a.e. for the treatment area per annual growing season. The total application to control floating and emerged weeds should not exceed 2 gallons formulation/acre per annual growing season.

## **Plants**

### *Selectivity*

Triclopyr TEA controls invasive species of aquatic macrophyte including Eurasian watermilfoil (*Myriophyllum spicatum*), parrotfeather (*Myriophyllum aquaticum*), waterhyacinth (*Eichhornia crassipes*), alligatorweed (*Alternanthera philoxeroides*), and purple loosestrife (*Lythrum salicaria*).

Triclopyr TEA does not control desirable native species like rushes (*Juncus* spp. and *Scirpus* spp.), cattails (*Typha* spp.), duckweed (*Lemna* spp.), Flatstem pondweed (*Potamogeton zosteriformis*), Coontail (*Ceratophyllum demersum*), Southern naiad (*Najas guadalupensis*), American pondweed (*Elodea canadensis* and water paspalum (*Paspalum fluitans*), and most species of algae including the green algae (*Spirogyra* spp., *Cladophora* spp., *Mougeotia* spp. *Volvox* spp., *Closterium* spp. and *Scenedesmus* spp.), *Chara* spp. and *Anabaena* spp. (Getsinger et al, 2000; Woodburn et al, 1993; Petty et al, 1998 and Green et al, 1989, Foster et al, 1997, Woodburn, 1988 and Houtman, 1997).

### *Non-target Aquatic Species*

Sensitive non-target aquatic species of plants are not likely to be affected at triclopyr concentrations of 2.5 ppm or less. At higher concentrations (2.5 ppm a.e.), southern naiad, American waterweed and coontail may be adversely impacted.

### *Algae*

Sensitive non-target aquatic species of algae are not likely to be affected at triclopyr concentrations of 2.5 ppm or less.

### *Endangered Plant Species*

Acute risk and endangered plant species levels of concern from runoff of triclopyr triethylamine salt during ground application are exceeded at  $\geq 9.0$  lb a.e./A (non-target plants inhabiting adjacent acreage) and  $\geq 1.5$  lb a.e./A (non-target plants inhabiting semi-aquatic areas) (Triclopyr RED). Aquatic use rates for this material are well below the use rates listed. Therefore, little or no harm to non-target terrestrial plants is expected due to either over-spray or the use of triethylamine salt treated irrigation water.

The total application of these products should not exceed 2.5 ppm a.e. for the treatment area per annual growing season.



### *Spray Drift*

Spray drift has the potential to damage sensitive terrestrial plants. Species of plant that appear to be especially susceptible are grapes, tobacco, vegetable crops and flowers or other desirable broadleaf plants. Even with low drift, onions and sunflowers may be adversely affected by rates of application typically used to control floating and emergent weeds (6 lbs a.e./acre) or wetland non-crop weeds (9 lbs a.e./acre). For example, treatment rates as low as 0.12 to 0.005 lbs a.e./acre may cause 25% damage to these sensitive crop species (EPA RED, 1998) (Table 9). These rates are exceeded even when drift is low (1% to 5%). Small amounts of drift can be an issue if many swaths are applied, and particularly if the product is applied from an aircraft (Forsythe et al, 1997).

### ***Permit Mitigations***

#### *Plants and algae*

Use as directed by the label.

#### *Endangered plant species*

- The total application of these products should not exceed 2.5 ppm a.e. for the treatment area per annual growing season.

### **Animals**

#### *Freshwater Invertebrates*

Triclopyr TEA and triclopyr acid are practically non-toxic to aquatic invertebrates and are not anticipated to be an acute or chronic risk due to their fairly short half-life (typically <5 days), low intrinsic toxicity to animals, and low tendency to accumulate in animal tissue.

While formulated triclopyr is not believed to be toxic to invertebrates, higher treatment rates (2.5 ppm a.e.) present a low to moderate risk.

Observed toxicity values for *Daphnia magna* (LC50 = 376 ppm a.e.), grass shrimp (LC50 = >234 ppm a.e.), pink shrimp (LC50 = 281 ppm a.e.), fiddler crab (>314 ppm a.e.) and crayfish (LC50 >103 ppm a.e.) place triclopyr TEA in the EPA's ecotoxicology categories of slightly toxic to practically non-toxic.

Other species of invertebrates are virtually unaffected by triclopyr TEA. For example, all other species of invertebrates that were tested have an LC50 of >100 ppm a.e.

In the field where triclopyr TEA was used to control Eurasian watermilfoil, waterhyacinth, or purple loosestrife, no invertebrate mortality or changes in invertebrate population structure was seen that could be attributed to the use of triclopyr TEA (Petty

et al, 1998, Green et al, 1989 and Gardner and Grue, 1996, Houtman et al, 1997, Foster et al, 1997 and Woodburn, 1988).

### *Amphibians*

No laboratory work was conducted on the effects of triclopyr TEA against amphibians. It is anticipated that amphibians will be affected by triclopyr TEA both acutely ( $LC_{50}$  = 82 to 182 ppm a.e. = 114 to 254 ppm a.i.) and chronically (MATC = 27 to 61 ppm a.e. = 38 to 93 ppm a.i.) at concentrations similar to that affecting fish. What little data is available from the field indicates that *Rana pipiens* adults and tadpoles remain common 11 weeks after treatment of the Columbia, Missouri pond site at rates of 2.5 ppm a.e. (Petty et al, 1998).

### *Avian/Birds*

Triclopyr acid is *slightly toxic* to birds when orally dosed or consumed in the diet. The triethylamine salt is *slightly toxic to practically non-toxic* when orally dosed or consumed in the diet. Reproduction of birds may be affected at levels greater than 100 ppm (RED).

Toxicity studies indicate that triclopyr and its products used as aquatic herbicides do not pose a significant acute or chronic risk to wild birds.

### *Fish, free-swimming aquatic invertebrates, and benthic invertebrates*

Most species of fish are tolerant of triclopyr TEA. Sensitive and environmentally relevant species such as the various salmon species (*Onchorhynchus* spp.) have demonstrated  $LC_{50}$ s that range between 96 and 182 ppm a.e. (Wan et al, 1987). These toxicity values place triclopyr TEA in the US EPA's ecotoxicological categories of slightly toxic ( $LC_{50}$  = >10 to 100 ppm) to practically non-toxic ( $LC_{50}$  = >100 ppm).

There have been no verified cases of toxicity to fish when triclopyr is used at the maximum use rate of 2.5 ppm a.e. When mortality occurs in the field after the use of triclopyr for the control of aquatic weeds, it is usually very low ( $\leq 11\%$ ) and attributable to an oxygen slump due to the presence of rapidly growing non-target aquatic plant species (Petty et al, 1998).

Triclopyr acid has been reported to be practically non-toxic to rainbow trout ( $LC_{50}$  = 117 ppm a.e. for rainbow trout) and bluegill sunfish (96-hour  $LC_{50}$  of 148 ppm a.e.) (Section 4, Tables 2, 17 and 18). Other authors have reported triclopyr acid to be moderately toxic with 96-hour  $LC_{50}$ s ranging from 5.3 ppm a.e. for pink salmon (*Oncorhynchus gorbuscha*) to 9.6 ppm a.e. for Chinook salmon (*Oncorhynchus tshawytscha*).

Triclopyr TEA is generally safe to fish, free-swimming aquatic invertebrates, and benthic invertebrates when the  $EC_{50}/LC_{50}$  is compared to typical four-day time-weighted average expected environmental concentration (TWA- EEC). However, when the toxicity of triclopyr is compared to other pesticides, it is classified according to the U.S. EPA Ecotoxicological Categories as slightly toxic ( $LC_{50}$  = >10 to 100 ppm) to embryo/larval and juvenile eastern oyster (*Crassostrea virginica*) rainbow trout (*Onchorhynchus*

*mykiss*), tidewater silverside (*Mendia beryllina*), chum salmon (*Onchorhynchus keta*) and fathead minnow (*Pimephales promelas*). However, triclopyr TEA is classified as practically non-toxic ( $LC_{50} > 100$  ppm) to bluegill sunfish (*Lepomis macrochirus*), other salmon species (*Onchorhynchus* spp.), *Daphnia magna*, grass shrimp (*Palaemonetes pugio*), pink shrimp (*Penaeus duorarum*), fiddler crab (*Uca pugiator*), and red swamp crayfish (*Procambarus clarki*). In general, triclopyr TEA can be considered to have very low toxicity to environmentally relevant fish and aquatic invertebrates. Triclopyr TEA appears to be extremely safe for use in the presence of threatened and endangered salmonid game-fish.

Triclopyr TEA appears to be safe for use in aquatic ecosystems. When comparing typical expected environmental concentrations (EEC) of triclopyr with laboratory  $LC_{50}$ s, the highest concentration that may be encountered immediately after application (2.5 ppm a.e. for control of submerged weeds or 4.4 ppm a.e. for control of floating and emerged weeds in shallow water) may affect more sensitive species. Fish and non-mollusk species would not be adversely impacted by these concentrations of triclopyr TEA. For example, the most sensitive fish species is rainbow trout with a 96-hour  $LC_{50}$  of 82 ppm a.e. and the most sensitive non-mollusk invertebrate is the red swamp crayfish with a 96-hour  $LC_{50}$  of  $>103$  ppm a.e. Since these species have  $LC_{50}$ s that are  $>10$ -fold greater than the EEC that occurs immediately after application, it is not likely that they would be adversely impacted by the effects of triclopyr TEA. However, the most sensitive mollusk is the embryo larval stage of the eastern oyster with a 48-hour  $EC_{50}$  for improperly developed embryo/larvae of 22 ppm a.e. Since the risk quotient generated from this  $LC_{50}$  and the lowest initial EEC is greater than the low level of concern (0.1), this segment of the biota may be harmed by exposure to triclopyr TEA. However, since the risk quotient is not higher than the high level of concern (0.5), this segment of the biota will probably not be adversely impacted if triclopyr is classified and used as a restricted use aquatic herbicide ( $RQ = EEC/EC_{50} = 4.4 \text{ ppm a.e.} / 22 \text{ ppm a.e.} = 0.2$ ). Some concern has been expressed that the eastern oyster is not an appropriate species to use in evaluations of risk for compounds that may not be used legally in estuaries. Furthermore, any concentration of triclopyr TEA entering an estuary would be greatly diluted by both untreated river/creek water and untreated sea water from the tidal action (CSI, Volume 5, Section 4, p. 63-64).

#### *Sea water challenge tests for salmon*

The following seawater challenge tests were done in support of the noxious weed control program at Ecology.

#### **Effects of Three Aquatic Herbicides on Smoltification in Juvenile Pacific Coho Salmon** by King, KA<sup>1</sup>, CE Grue<sup>1</sup>, JM Grassley<sup>1</sup>, CA Curran<sup>1</sup>, WW Dickhoff<sup>2</sup>, and JA Winton<sup>3</sup>

Herbicides are frequently used to control exotic or nuisance aquatic plants. Utilization of herbicides in Integrated Pest Management (IPM) plans to control aquatic weeds has been hampered by court injunctions directed at the non-target

toxicity of active herbicidal ingredients. Unfortunately, adequate data on the non-target toxicity of aquatic herbicides to aquatic resources are lacking, thereby threatening the permitting process and the success of IPM strategies. Recent declines in several species/stocks of salmon and the emphasis of management and regulatory agencies to restore these populations heighten concerns. Our objective was to determine if label application rates for three commonly used aquatic herbicides impair smoltification in juvenile Pacific salmon, using coho (*Oncorhynchus kisutch*) as a model. The herbicides and water concentrations selected for study were Sonar®PR (active ingredient: fluridone; 10, 90 ppb), REWARD® (active ingredient: diquat dibromide; 0.34, 1.37 ppm), and RENOVATE® (active ingredient: triclopyr; 0.75, 2.50 ppm). Fish (mean = 22g; 20 fish/tank) were exposed to the herbicides or negative control (4 tanks/treatment) for 96 h under static conditions (11 C) and then transferred directly into flowing seawater (salinity = 27 ppt; 10 C) for 14 d. Five fish per tank were sacrificed after exposure to the chemicals, and after 1, 7 and 14 d in seawater. Endpoints were survival, body weight and fork length, muscle water content, hepatosomatic index, plasma sodium and chloride concentrations, gill ATPase, and gill histology. Tests for each herbicide were conducted concurrently. Actual concentrations were similar to nominal with the exception of fluridone (1, 10 ppb) due to a calculation error. All fish survived the chemical exposures and the first 7 d in seawater. Two fish exposed to the low concentration of REWARD died during the second week in seawater. The fish were from the same tank; no mortality was observed in other three replicate tanks. Necropsies of the two fish did not reveal any gross anomalies. Statistically significant decreases in plasma Cl concentrations were detected in fish exposed to the low and high levels of RENOVATE and REWARD while in freshwater compared to controls. Significant decreases in plasma Na and Cl were observed in REWARD-exposed fish after 24 h in seawater. Effects were short-lived and plasma ion concentrations were similar among treated and control fish after 7 and 14 d in seawater. Hepatosomatic index and muscle water content did not differ between treated fish and controls in either fresh or seawater. A few differences were detected in gill ATPase between treated and control fish, but effects were not consistent. Data on gill histology have yet to be analyzed statistically. Preliminary results suggest that, at the chemical and seawater exposures tested, the herbicides are unlikely to affect seawater adaptation in free-living juvenile Pacific salmon.

<sup>1</sup>Washington Cooperative Fish and Wildlife Research Unit, School of Aquatic and Fishery Sciences, Box 355202, University of Washington, Seattle, WA 98195;

<sup>2</sup>National Marine Fisheries Service - Montlake Laboratory, Seattle, WA 98112;

<sup>3</sup>US Geological Survey, Biological Resources Division, Western Fisheries Research Center, Seattle, WA 98115.

### *Mammals*

There are two common routes of exposure of livestock and terrestrial wildlife to aquatic applications of Renovate®. The two routes are exposure through drinking water treated

with products containing triclopyr or eating aquatic plants, fish, or other aquatic organisms from the treatment site. Based on acute and chronic studies, triclopyr and its products used as aquatic herbicides do not pose a significant acute or chronic risk to terrestrial mammals.

### *Threatened and Endangered Species*

Minimal effects to threatened and endangered species are expected from application of aquatic herbicides containing triclopyr. Mitigation of possible effects on listed endangered species is best accomplished by following the mitigation sections for terrestrial plants, birds and animals. As stated previously, the best way to mitigate possible effects on all terrestrial species is to follow the directions listed on the label.

Other mitigation measures involve the contact of WDFW by the issuer of the permit to ascertain if any endangered species may be affected by the application of the chemical to the water body in question. Questions asked by the permit granter would ascertain if any resident endangered bird or animal species are known to use the water body in question (or its shorelines or islands) as breeding or forage areas, or if the application coincides with the migration of any endangered species. If endangered species are present, mitigation measures may involve postponing application until after the breeding season or postponement of application until after migration of the species in question. Use of an alternate means of control (i.e. mechanical) may also be an option if the risk is determined to be too great to the species in question.

### ***Permit Mitigation***

#### *Invertebrate biota*

#### *Wetland species*

#### *Fish*

Use as directed by the label.

#### *Avian*

- If the chemicals are applied according to the label, the effect on terrestrial wildlife should be minimal. Even though triclopyr products used as aquatic herbicides do not pose a significant risk to terrestrial wildlife, the following measures should be considered prior to all aquatic herbicide applications. One possible mitigation measure would be not allowing applications if large populations of birds use shorelines or islands in the water body to be treated for nesting until after nesting is complete. Another mitigation measure would be to time applications to avoid migratory waterfowl and other bird species that use certain water bodies during migration. Efforts to avoid effects on migratory and nesting birds would best be coordinated with The Washington State Department of Fish and Wildlife (WDFW).

## Water, Land and Shoreline Use

### *Public Water Supplies - Potable Water*

The Reference Dose (RfD), the amount of triclopyr residuals that could be consumed daily over a lifetime without adverse effects, was established at 0.05 mg/kg/day, based on the two-generation reproduction toxicity study in rats with a NOEL (no observed effect level) of 5.0 mg/kg/day, the lowest dose tested. At the next dose level (25 mg/kg/day), an increased incidence of proximal tubular degeneration of the kidneys was observed in P1 and P2 parental rats in this study (EPA R.E.D. Facts, 1998).

Triclopyr and its toxic metabolite TCP degrade and dissipate rapidly through chemical, biological, and physical processes (Various authors in Houtman et al, 1997). Concentrations of triclopyr in sites with short half-lives will typically fall below the temporary drinking water tolerance within one to three days of application. In areas with short triclopyr half-lives, the metabolite TCP is often not detected after the day of treatment, but has been detected at concentrations of 0.05 to 0.14 ppm in Lake Seminole, Georgia (CSI, Volume 5, Sect. 3, p. 40).

### *Potable Water*

It has also been proposed as part of the tolerance petition, that the potable water setback be 0.25 miles in order to ensure residue levels remain below 0.5 ppm (proposed allowable drinking water tolerance). This set back distance was based on the results of several field dissipation studies (Woodburn, 1988 Houtman et al, 1997, Foster et al, 1997). However, recent modeling work (Ritter and Peacock, 2000) indicates that the setback distance should vary with the concentration used and the number of acres treated. At the maximum use rate (2.5 ppm) used to treat >16 acres, the setback distance from potable water intakes should be at least 2000 feet.

### *Ground Water*

Highly mobile and water soluble compounds are more likely to reach ground water. Triclopyr is highly mobile ( $K_d = 0.165$  to  $0.975$ ), and highly water soluble.

While triclopyr exceeds the mobility and persistence triggers used to recommend restricted use, triclopyr does not meet detection triggers for recommending restricted use due to limited monitoring data (Hoheisel et al, 1992 in EPA RED, 1998).

In one EPA study (EPA, 1992), three hundred seventy-nine wells were sampled for triclopyr, and only five detections of triclopyr residues in ground water were reported. All detections were far below levels of concern. The maximum concentration reported was 0.58 ppb.

Public water supply systems are not required to sample for triclopyr, as triclopyr is currently not regulated under the Safe Drinking Water Act (SDWA). There is no

maximum contaminant level (MCL) or Drinking Water Lifetime Health Advisory (HAL) for triclopyr. However, there is a proposed MCL of 500 ppb and an estimated HAL of 350 ppb.

Due to the limited amount of data collected, it is difficult to determine if triclopyr will have an adverse impact on sensitive well recharge areas. Although EPA does not currently have surface or ground water advisories on triclopyr, surface and ground water studies may be necessary to determine the potential for triclopyr to leach under its new aquatic use labeling (EPA RED, 1998; Getsinger et al, 1997; Green et al, 1989; Getsinger et al, 2000; Petty et al, 1998 and Petty et al 1998).

### *Swimming*

The only health concerns from triclopyr for swimming are minor eye irritation and exposure to children immediately after application. The risk of eye irritation and overexposure for children decreases rapidly because of dilution. A mandatory waiting time after application before swimming is allowed mitigates the risk.

Exposure and risk calculations were determined for hypothetical situations involving ingestion and dermal contact with treated water while swimming and drinking potable water. Calculation of triclopyr exposures utilized the swimmer's weight, the skin surface area available for exposure, the amount of time spent in the treated water containing 2.5 and 0.5 ppm triclopyr, amount of water swallowed while swimming over specific time periods, and the estimated human skin permeability coefficient.

Risk analyses were completed for various populations. The most sensitive population was found to be children who swim for three hours and ingest water while swimming. However, a child would have to ingest 3.5 gallons of lake water where triclopyr had been recently applied to cause risk factors to be exceeded.

Based on the label use directions and the results of the triclopyr toxicology studies, the aggregate or combined daily exposure to the chemical from aquatic herbicidal weed control does not pose an adverse health concern.

The Washington State Dept. of Health has recommended a 12-hour restriction for re-entry into triclopyr treated water to assure that the eye irritation potential and any other adverse effects will not occur. WDOH also recommends that those wanting to avoid even small exposures can wait one to two weeks following application when the triclopyr residues have dissipated from the water and sediments (WDOH, 1999).

### *Drift*

The main methods of using Renovate® and Garlon® 3A largely preclude the effects of drift. This liquid product is either injected by subsurface methods (which precludes drift) or applied as large droplets at low pressure which mitigates the effects of drift. It is also recommended that a thickening agent be used to control drift when applying liquid herbicides to the water surface or to wetland associated weeds.

The Garlon® 3A proposed label states the following: “Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants are near enough to be injured. It is suggested that a continuous smoke column at or near the spray site or a smoke generator on the spray equipment be used to detect air movement, lapse conditions, or temperature inversions (stable air). If the smoke layers or indicates a potential of hazardous spray drift, do not spray.” Spray pressures should be kept low enough to provide coarse droplets. The spray boom should be kept as close to the ground or water surface as possible. In addition, a thickening agent or a high viscosity inverting system should be used to prevent drift.

### ***Permit Mitigation***

#### *Water intakes and Drinking water*

Follow label instructions.

#### *Irrigation*

Follow label directions

#### *Ground Water*

- Use according to label directions. Ground water or sediment monitoring is required prior to any third application of triclopyr on a previously treated site planned within a three-year period. Evidence of persistence of triclopyr or TCP in sediment or ground water is basis for denial of the third application.

#### *Swimming*

- The Washington State Dept. of Health has recommended a 12-hour restriction for re-entry into triclopyr treated water to assure that the eye irritation potential and any other adverse effects will not occur. WDOH also recommends that those wanting to avoid even small exposures can wait 1-2 weeks following application when the triclopyr residues have dissipated from the water and sediments (WDOH, 1999).

#### *Fish Consumption*

#### *Agriculture*

#### *Irrigation*

Follow label directions.



### *Data Gaps and Considerations*

Since triclopyr bioaccumulates at low levels (~1.0 to 2.0 in crayfish and clams), further evaluation of the accumulation effects of triclopyr on clams and crayfish should be considered before establishing residue tolerance limits on these species. The current proposed residue tolerance for fish and shellfish is 0.2 ppm.

Wetland (forestry) herbicides may be of particular concern to Native Americans. Forestry products are harvested by Native Americans and are used in their diets, in the making of traditional basketry, for medicinal purposes and ceremonial activities. Work is currently being conducted to determine if these exposure scenarios may affect Native Americans in a manner not reflected in the current assessment.

No laboratory work was conducted on the effects of triclopyr TEA against amphibians. It is anticipated that amphibians will be affected by triclopyr TEA both acutely ( $LC_{50} = 82$  to 182 ppm a.e. = 114 to 254 ppm a.i.) and chronically (MATC = 27 to 61 ppm a.e. = 38 to 93 ppm a.i.) at concentrations similar to that affecting fish. What little data is available from the field indicates that *Rana pipiens* adults and tadpoles remain common 11 weeks after treatment of the Columbia, Missouri pond site at rates of 2.5 ppm a.e. (Petty et al, 1998).

## **4. Mitigation Summary for Triclopyr TEA**

<b>Conditions of Treatment</b>	<b>Mitigation</b>
Drift	Follow label directions.
Odor	Aerial applications are not allowed so this should not be an issue.
Inhalation	Persons involved with mixing or applying should follow the directions on the label for safety. Application should be by direct injection under the water and not applied by aerial or spray application.
Irrigation	Follow label directions.
Surface water/fish and other aquatic life	Follow label directions.
Wetlands, estuaries and marshes that treated water may flow into.	The total application of these products should not exceed 2.5 ppm a.e. for the treatment area per annual growing season. The total application of these products to control floating and emerged weeds should not exceed 2 gallons formulation/acre per annual growing season.
Non-target plants, including endangered plant species	The total application of these products should not exceed 2.5 ppm a.e. for the treatment area per annual growing season.
Invertebrates	If triclopyr TEA is treated as a restricted use herbicide, it should not cause adverse impact to the invertebrate biota.

Birds	Do not apply when large populations of birds use shorelines or islands in the water body to be treated for nesting until after nesting is complete. Avoid migratory waterfowl and other bird species that use certain water bodies during migration.
Threatened and endangered species	<p>Mitigation of possible effects on listed endangered species is best accomplished by following the mitigation sections for terrestrial plants, birds and animals.</p> <p>Follow the directions listed on the label.</p> <p>Other mitigation measures involve the contact of WDFW by the issuer of the permit to ascertain if any endangered species may be affected by the application of the chemical to the water body in question.</p>
Swimming	The Washington State Dept. of Health has recommended a 12-hour restriction for re-entry into triclopyr treated water to assure that the eye irritation potential and any other adverse effects will not occur. WDOH also recommends that those wanting to avoid even small exposures can wait 1-2 weeks following application when the triclopyr residues have dissipated from the water and sediments (WDOH, 1999).
Potable Water	Follow label directions.
Fishing/Fish consumption/shellfish consumption	Follow label directions.
Ground water	Use according to label directions. Ground water or sediment monitoring is required prior to any third application of triclopyr on a previously treated site planned within a three-year period. Evidence of persistence of triclopyr or TCP in sediment or ground water is basis for denial of the third application.

## References

The primary reference for this document is the Supplemental Environment Impact Statement Assessments of Aquatic Herbicides: Study No. 00713, Volume 5, Triclopyr, by Compliance Services International, Tacoma, Washington, submitted to the Dept. of Ecology under a grant contract. The five sections are as follows:

- Section 1 – Label Description & History, 65 pp.
- Section 2 – Chemical Characteristics, 10 pp.
- Section 3 – Environmental Fate, 94 pp.
- Section 4 – Environmental Effects, 160 pp.
- Section 5 – Human Health Effects, 47 pp.

The following are references cited in the above document:

### Section 1 – Label Description & History

1. Brian Database. 1999. Online EPA Database that Summarizes Ecotoxicological Data that EPA Uses for Ecotoxicological Assessments.
2. Ebasco. 1993. Final Report, Element E. Chemical Methods Only: Environmental Effects of Glyphosate and 2,4-D, for Washington State Department of Ecology.
3. Ecology. 1980. Environmental Impact Statement. Aquatic Plant Management. DRAFT. February 1980.
4. Ecology. 1989. Draft Environmental Impact Statement Supplement. State of Washington Aquatic Plant Management Program. U.S. Army Corps of Engineers, Seattle District.
5. Ecology. 1991 & 1992. Aquatic Plants Management Program for Washington State. Final Supplemental Impact Statement and Responsiveness Summary Volumes 1 & 2.
6. ECOTOX Database. 1999. Online EPA Database that Summarizes Ecotoxicological Collected from Reviewed Literature as well as the Un-reviewed Brian Database.
7. EPA Registration Eligibility Decision – Triclopyr. EPA 738-R-98-011.
8. Houtman, B.A.; Foster, D.R.; Getsinger, K.D. and Petty D.G., 1997. Aquatic Dissipation in Lake Minnetonka, DowElanco, ENV94011. MRID 44456102.
9. Foster, D.R.; Getsinger, K.D. and Petty D.G., 1997. Aquatic Dissipation of triclopyr in a Whole Pond Treatment, DowElanco, ENV94012. MRID 44456103.
10. Gardner, S.C. and Grue, C.E., 1996. Effects of Rodeo and Garlon® 3A on Nontarget Wetland Species in Central Washington. Environmental Toxicology and Chemistry 15(4): 441-451.

12. 11. Getsinger, K.D.; Petty, D.G.; Madsen, J.D.; Skogerboe, J.G.; Houtman, B.A.; Haller, W.T. and Fox, A.M., 2000. Aquatic Dissipation of Herbicide Triclopyr in Lake Minnetonka, Minnesota. *Pest Manag. Sci.* 56: 388-400.
13. Garlon® 3A -- Proposed Label. Specimen Label from DowElanco, 2000.
14. Getsinger, K. 1999, Personal Communication.
14. Green, W.R.; Westerdahl, H.E.; Joyce, J.C. and Haller, W.T., 1989. Triclopyr (Garlon® 3A) Dissipation in Lake Seminole, Georgia. Miscellaneous Paper A-89-2, US Army Engineer Waterways Experiment Station Vicksburg, MS.
15. Petty, D.G.; Jestig, K.D.; Madsen, J.D. Skogerboe, J.G.; Haller, W.T. and Fox, A.M.; Houtman, B.A., 1998. Aquatic Dissipation of Herbicide Triclopyr in Lake Minnetonka, Minnesota. U.S. Army Corp of Engineers. Technical Report A-98-1.
16. Renovate® -- Experimental Use Label, 1999. Experimental Use Permit No 62719-EUP-1.
17. Resource Management, Inc., 1999. Report on 1999 Treatment of Lake Steilacoom.
18. Ritter, A.M. and Peacock, A.L., 2000. Aquatic Dissipation Modeling of Triclopyr. Waterborne Environmental, Inc. Dow Agro Sciences, LLC WEI 396.06. LLC 30398118-5023-1. MRID 45047901.
19. Wan, M.T.; Moul, D.J. and Watts, R.G., 1987. Acute Toxicity to Juvenile Pacific Salmonids of Garlon 3A™, Garlon 4™, triclopyr ester and their Transformation products: 3,5,6 trichloro-2-pyridinol (TCP) and 2-methoxy-3,5,6-trichloropyridine (TMP). *Bulletin of Environmental Contamination and Toxicology* 39: 721-728.
20. Westerdahl, Howard E., and K.D. Getsinger., eds. 1988. "Aquatic Plant Identification and Herbicide Use Guide; Volume II: Aquatic Plants and Susceptibility to Herbicides," Technical Report A-88-9, US Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
21. Woodburn, K.B., 1988. The Aquatic Dissipation of Triclopyr in Lake Seminole Georgia. Dow Chemical. GH-C 2093. MRID 41714304.
22. Woodburn, K.B.; Green, W.R. and Westerdahl, H.E., 1993. Aquatic Dissipation of Triclopyr in Lake Seminole, Georgia. *J. Agricultural and Food Chemistry* 41: 2172- 2177.
23. Washington State Department of Ecology Permits for Renovate® Specialty Herbicide Applications, 1998 to 2000.

## **Section 2 – Chemical Characteristics**

1. EPA. 1998. Reregistration Eligibility Decision (RED): Triclopyr. United States Environmental Protection Agency. Washington, D.C.

2. Hall, S. 1999. "Group B-Physical/Chemical Properties for Garlon 3A, a Liquid End-Use Product Containing Triclopyr TEA". Unpublished Study prepared by Dow AgroSciences LLC. Indianapolis, Indiana.
3. Purdue University. 2000. Herbicide Mode-Of-Action Summary. Cooperative Extension Service, West Lafayette, IN. [www.agcom.purdue.edu/AgCom/Pubs/WS/WS-23.htm](http://www.agcom.purdue.edu/AgCom/Pubs/WS/WS-23.htm). Accessed June 26, 2000.
4. SePRO. 2000. "Renovate Aquatic Herbicide Technical Bulletin". SePRO Corporation, Carmel, Indiana.

### Section 3 – Environmental Fate

1. Abdelghani, A.A. 1995. Assessment of exposure of workers applying herbicide mixtures, toxicity and fate of these mixtures in the environment State Project #736-14-0067. Summary Report. Louisiana Department of Transportation and Development. Note: more complete data cited in Obenshain et al., 1997.
2. Bidlack, H.D. (no date). Dow Chemical. Unpublished report. Cited in McCall and Gavit, 1986.
3. Bollag, J. M., and S.-Y. Liu. 1990. Biological transformation processes of pesticides. In: Pesticides in the Soil Environment. SSSA Book Series, No. 2, pp. 169-211. Soil Science Society of America, Madison, Wisconsin.
4. Branham, B.E. 1994. Herbicide fate in turf. Pages 109-151 *in* Turgeon, A.J. (ed.), Turf weeds and their control, American Society of Agronomy, Madison, Wisconsin.
5. Chapman, R.A. and C.M. Cole. 1982. Observations on the influence of water and soil pH on the persistence of insecticides. *Journal of Environmental Science and Health*, B17: 487-504.
6. Cleveland, C.B., and D.L. Holbrook. 1991. A hydrolysis study of triclopyr. Dow AgroSciences, Indianapolis, Indiana. Cited in Getsinger et al, 2000.
7. Dilling, W.L. et al. 1984a. Unpublished data of The Dow Chemical Company. Cited in Woodburn et al., 1993b.
8. Dilling, W.L., L.C. Lickly, T.D. Lickly, P.G. Murphy, and R.L. McKellar. 1984b. Organic Photochemistry. 19. Quantum yields for O,O-diethyl-0-(3,5,6-trichloro-2-pyridinyl) phosphorothioate (chloropyrifos) and 3,5,6-trichloro-2-pyridinol in dilute aqueous solutions and their environmental phototransformation rates. *Environmental Science and Technology*, 18:540-543.
9. Ebasco. 1993. Chemical Methods Only: Environmental Effects of Glyphosate and 2,4-D. Washington Department of the Ecology Final Report Element E.
10. Feng, Y., R.D. Minard, and J.M. Bollag. 1998. Photolytic and microbial degradation of 3,5,6-trichloro-2-pyridinol. *Environmental Toxicology and Chemistry*, 17(5): 814-819.
11. Feng, Y., K.D. Racke, and J.M. Bollag. 1997. Isolation and characterization of a chlorinated-pyridinol-degrading bacterium. *Applied Environmental Microbiology*, 63: 4096-4098.

12. Gardner, S.C., and C.E. Grue. 1996. Effects of Rodeo (R) and Garlon (R) 3A on nontarget wetland species in central Washington. *Environmental Toxicology and Chemistry*, 14(4): 441-451.
13. Garlon® 3A Label. Proposed in 2000.
14. Getsinger, K.D., D.G. Petty, J.D. Madsen, J.G. Skogerboe, B.A. Houtman, W.T. Haller, and A.M. Fox. 2000. Aquatic dissipation of the herbicide triclopyr in Lake Minnetonka, Minnesota. *Pest Management Science*, 56: 388-400.
15. Getsinger, K.D., E.G. Turner, J.D. Madsen, M.D. Netherland. 1997. Restoring native vegetation in a Eurasian watermilfoil dominated plant community using the herbicide triclopyr. *Regulated Rivers - Research and Management*, 13(4): 357-375.
16. Getsinger, K.D. and H.E. Westerdahl. 1984. Field evaluation of Garlon 3A (triclopyr) and 14-ACE-B (2,4-D BEE) for the control of Eurasian milfoil. *Miscellaneous Paper A-84-5*. US Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
17. Green, W.R., H.E. Westerdahl, J.C. Joyce, and W.T. Haller. 1989. Triclopyr (Garlon 3A) dissipation in Lake Seminole, Georgia. *Miscellaneous Paper A-89-2*, US Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.
18. Hamaker, (no date). Dow Chemical. Personal communication cited in McCall and Gavit, 1986.
19. Hamaker, J.W. 1975. The hydrolysis of triclopyr in buffered distilled water. Dow AgroSciences, Indianapolis, Indiana. Cited in Getsinger et al., 2000.
20. Houtman, B.A., D.R. Foster, K.D. Getsinger, and D.G. Petty. 1997. Aquatic dissipation of triclopyr in Lake Minnetonka, Minnesota. *GH-C 4449*. Dow AgroSciences, Indianapolis, Indiana. Cited in Petty et al., 1998.
21. Johnson, W.G., and T.L. Lavy. 1992. Dissipation of 2,4-D and triclopyr under controlled temperature and water potential. *Proceedings of the Southern Weed Science Society*, 45: 366.
22. Johnson, W.G., T.L. Lavy, and E.E. Gbur. 1995a. Persistence of triclopyr and 2,4-D in flooded and nonflooded soils. *Journal of Environmental Quality*, 24: 493-497.
23. Johnson, W.G., T.L. Lavy, and E.E. Gbur. 1995b. Sorption, mobility and degradation of triclopyr and 2,4-D on four soils. *Weed Science*, 43: 678-684.
24. Jotcham, J.R., D.W. Smith, and G.R. Stephenson. 1989. Comparative persistence and mobility of pyridine and phenoxy herbicides in soil. *Weed Technology*, 3: 155-161.
25. Larson, A.G. 1996. Washington State Pesticide Monitoring Program. Pesticides in Washington State Ground Water – A Summary Report, 1988-1995. Pesticides in Ground Water No. 9. Washington State Department of Ecology. Publication No. 96-303.
26. Lee, C.H., P.C. Oloffs, and S.Y. Szeto. 1986. Persistence, degradation and movement of triclopyr and its ethylene glycol butyl ether ester in a forest soil. *Journal of Agricultural and Food Chemistry*, 34: 1075-1079.

27. Leng, M.L., E.M.K. Leovey, and Zubkoff. 1995. *Agrochemical Environmental Fate: State of the art*. CRC Press. Lewis Publishers, Boca Raton. 410 pages
28. McCall, P.J. and P.D. Gavit. 1986. Aqueous photolysis of triclopyr and its butoxyethyl ester and calculated environmental photodecomposition rates. *Environmental Toxicology and Chemistry*, 5: 879-886.
29. McCall, P.J., D.A. Laskowski, and H.D. Bidlack. 1988. Simulation of the aquatic fate of triclopyr butoxyethyl ester and its predicted effects on Coho salmon. *Environmental Toxicology and Chemistry*, 7(7): 517-528.
30. Miller, G.C. and R.G. Zepp. 1983. Extrapolating photolysis rates from the laboratory to the environment. *Residue Rev*, 85: 89-110.
31. Newbold, C., 1975. Herbicides in Aquatic Systems. *Biological Conservation* 7(2): 97.
32. Norris, L.A., M.L. Montgomery, and L.E. Warren. 1987. Triclopyr persistence in western Oregon hill pastures. *Bulletin of Environmental Contamination and Toxicology*, 39:134-141.
33. Obenshain, K.R., M.C. Metcalf, A.A. Abdelghani, J.L. Regens, D.G. Hodges, and C.M. Swalm. 1997. Spatial analysis of herbicide decay rates in Louisiana. *Environmental Monitoring and Assessment*, 48: 307-316.
34. Petty, D.G., K.D. Getsinger, J.D. Madsen, J.G. Skogerboe, W.T. Haller, A.M. Fox and B.A. Houtman. 1998a. The aquatic dissipation of triclopyr in Lake Minnetonka, Minnesota. Technical Report A-98-01, U.S Army Engineer Waterways Experiment Station, Vicksburg, Mississippi. 96 pages.
35. Petty, D.G., J.G. Skogerboe, K.D. Getsinger, D.R. Foster, and J.W. Fairchild. 1998b. Aquatic dissipation of triclopyr in a whole-pond treatment (final report). Technical Report A-98-06, U.S Army Engineer Waterways Experiment Station, Vicksburg, Mississippi. 133 pages.
36. Racke, K.D., J.R. Coats, and K.R. Titus. 1988. Degradation of chlorpyrifos and its hydrolysis product, 3,5,6-trichloro-2-pyridinol, in soil. *Journal of Environmental Science and Health B*, 23: 527-539.
37. Reeves, G.L. 1995. The mobility of triclopyr under laboratory and field conditions. Pages 223-229 in Walker, A. et al., *Pesticide movement to water: Proceedings of a symposium held at the University of Warwick, Coventry, UK, on 3-5 April 1995*. Monograph No. 62, British Crop Protection Council, Farnham, UK.
38. Renovate® EUP, 1997. Federal Experimental Use Permit, November 25, 1997.
39. Renovate® Experimental Label. L1A/Renovate/EUP Renewal/02-15-99
40. Ritter, A.M. and Peacock, A.L. 2000. Aquatic Dissipation of Triclopyr. Dow AgroSciences. LLC 30398118-5023-1. MRID 45047901.
41. Solomon, K.R., C.S. Bowhey, K. Liber, and G.R. Stephenson. 1988. Persistence of hexazinone (Velpar), triclopyr (Garlon), and 2,4-D in a northern Ontario aquatic environment. *Journal of Agricultural and Food Chemistry*, 36: 1314-1318.

42. Stephenson, G.R., K.R. Solomon, C.S. Bowhey, and K. Liber. 1990. Persistence, Leachability, and lateral movement of triclopyr (Garlon) in selected Canadian forest soils. *Journal of Agricultural and Food Chemistry*, 38: 584-588.
43. Szeto, S.Y. 1993. Determination of kinetics of hydrolysis by high-pressure liquid chromatography: Application to hydrolysis of the ethylene glycol butyl ester of triclopyr. *Journal of Agricultural and Food Chemistry*, 41(7): 1118-1121.
44. Thompson, D.G., B. Staznik, D.D. Fontaine, T. Mackey, G.R. Oliver, and J. Troth. 1991. Fate of triclopyr ester (release) in a boreal forest stream. *Environmental Toxicology and Chemistry*, 10(5): 619-632.
45. USEPA (US Environmental Protection Agency). October, 1998. Reregistration Eligibility Decision (RED) - Triclopyr. EPA document 738-R-98-011. 213 pages.
46. USGS. 1997. Pesticides in selected Small Streams in the Puget Sound Basin, 1987 to 1995. USGS Fact Sheet 067-97.
47. USGS. 1999. Pesticides Detected in Urban Streams During Rainstorms and Relations to Retail Sales of Pesticides in King County, Washington. USGS Fact Sheet 097-99.
48. Wan, M.T.; D.J. Moul and R.G. Watts. 1987. Acute Toxicity to Juvenile Pacific Salmonids of Garlon 3A™, Garlon 4™, Triclopyr, Triclopyr Ester and Their Transformation Products: 3,5,6-Trichloro-2 Pyridinol and 2-Methoxy-3,5,6-Tricholopyridine. *Bulletin of Environmental Contamination and Toxicology* 39: 721-728.
49. Wauchope, R.D., T.M. Butler, A.G. Hornsby, P.W.M. Augustyn-Beckers, and J.P. Burt. 1992. The SCS/ARS/CES pesticide properties database for environmental decision-making. *Environmental Contamination and Toxicology*, 123:1-64.
50. Woodburn, K.B. 1988. The Aquatic Dissipation of Triclopyr in Lake Seminole. Dow Chemical. GH-C 2093 MRID 41714304.
51. Woodburn, K.B., and W. Cranor. 1987. Aerobic aquatic metabolism of [14C] triclopyr. Dow AgroSciences, Indianapolis, Indiana. Cited in Getsinger et al., 2000.
52. Woodburn, K.B., R.R. Batzer, F.R. White, and M.R. Schultz. 1990. The aqueous photolysis of triclopyr. Dow AgroSciences, Indianapolis, Indiana. Cited in Getsinger et al, 2000.
53. Woodburn, K.B., F.R. Batzer, F.H. White, M.R. Schultz. 1993a. The aqueous photolysis of triclopyr. *Environmental Toxicology and Chemistry*, 12: 43-55.
54. Woodburn, K.B.; D.D. Fontaine and H.F. Richards. 1988. A Soil Adsorption/Desorption Study of Triclopyr. Dow Chemical. GHC-2017. MRID 40749801.
55. Woodburn, K.B., W.R. Green, and H.E. Westerdahl. 1993b. Aquatic dissipation of triclopyr in Lake Seminole, Georgia. *Journal of Agricultural and Food Chemistry*, 41: 2172-2177.



#### Section 4 – Environmental Effects

1. Aislabie, J and Lloyd-Jones. 1995. A Review of Bacterial Degradation of Pesticides. Australian Journal of Soil Research 33(6): 925-942.
2. Bailey, R.E. and D.L Hopkins. 1987. ((3,5,6-trichloro-2-pyridinyl)oxy) acetic acid, triethylamine Salt – Octanol/Water partition coefficient. Dow Chemical Company. ES-DR-0114-8261-2. MRID 4121910.
3. Barron, M.G., S.C. Hansen and T. Ball. 1991. Pharmacokinetics and Metabolism of Triclopyr in Crayfish (*Procambarus clarki*). Drug Metabolism and Disposition. 19(1): 163-167.
4. Batchelder, T.L. 1973. Acute Toxicity of DOWCO 233 (3,5,6-trichloro-2pyridyloxyacetic) Acid and Two Derivative. Dow Chemical. WDL 7305. MRID 00049637.
5. Bauriedel, W.R. 1983. The Metabolic Fate of 14C-Triclopyr Fed to Lactating Goat. Dow Chemical. #11562.
6. Berril, M., S. Bertram, L. McGillvray, M. Kolohon, and B. Pauli. 1994. Effects of Low Concentrations of Forest –use Pesticides on Frog Embryos and Tadpoles. Environmental Toxicology and Chemistry 13(4): 657-664.
7. Boca Raton, FL. In Washington State, 1993.
8. Brian Database. 1999. EPA Database for Ecotoxicological Effects.
9. Campbell K.R., S.M. Bartell and J.L.Shaw. 2000. Characterizing Aquatic Ecological Risks from Pesticides Using A Diquat Dibromide Case Study. II. Approaches Using Quotients and Distributions. Environmental Toxicology and Chemistry 19(3): 760-774.
10. Cowgill, U.M. and D.P. Milazzo. 1987. Triclopyr Triethylamine Salt: The Fourteen-Day Toxicity to *Lemna gibba* L. G-3 (Duckweed). Dow Chemical Company. ES-DR-0287-0871-1. MRID 41633709.
11. Cowgill, U.M., D.P. Milazzo and B.D.Landenberger. 1989. Toxicity of Nine Benchmark Chemicals to *Skeletonema costatum* a Marine Diatom. Environmental Toxicology and Chemistry 8: 451-455.
12. Cowgill, U.M., D.P. Milazzo and B.D. Landenberger. 1988. A Comparison of the Effect of Triclopyr Triethylamine Salt (Garlon® 3a) on Two Species of Duckweed Examined for Sensitivity During a Fourteen Day Period. Dow Chemical Company. ES-DE-0003-07070-2.
13. Dingledine, J. 1985. Triclopyr Acid. An Acute Contact Toxicity Study with Honey Bees. Final Report. Wildlife International, Ltd. Lab. Project 103-239. MRID 40356602.
14. EBASCO Environmental. 1993. Chemical Methods Only: Environmental Effects Glyphosate and Diquat. Environmental Impact Statement Prepared for the Washington State Department of Ecology, Element E.
15. Ecology. 1980. Environmental Impact Statement. Aquatic Plant Management Draft. February, 1980.
16. Ecology. 1989. State of Washington Aquatic Plant Management Program. Draft Environmental Impact Statement.
17. Ecology. 1991/1992. Aquatic Plants Management Program for Washington State. Final Supplemental Environmental Impact Statement and Responsiveness Summary. Volume 1.

18. Ecology. 1991/1992. Aquatic Plants Management Program for Washington State. Draft Supplemental Environmental Impact Statement and Responsiveness Summary. Volume 2: Appendices.
19. Ecology. 2000. Supplemental Environmental Impact Statement. Assessment of Aquatic Herbicides. Appendix 1. Volume 3: 2,4-D. Washington State Department of Ecology. Publication Number 00-10-043.
20. El-Deen, M.A.S. and W.A. Rogers. 1992. Acute Toxicity and Some Hematological Changes in Grass Carp Exposed to Diquat. *Journal of Aquatic Animal Health* 4: 277-280.
21. EPA. 1982. Pesticide Assessment Guidelines. Subdivision E. Hazard Evaluation: Wildlife and Aquatic Organisms. EPA 540/9-82-024. PB-153908.
22. EPA RED. 1998. Registration Eligibility Decision – Triclopyr. EPA 738-R-98-01. October, 1998.
23. Foster, D.R., K.D. Getsinger, and D.G. Petty. 1997. Aquatic Dissipation of Triclopyr in Lake Minnetonka, DowElanco, ENV94011. MRID 44456102.
24. Forsythe, D.J., P.A. Martin and G.G. Shaw. 1997. Effects of Herbicides on Two Submersed Aquatic Macrophytes, *Potamogeton pectinatus* L. and *Myriophyllum sibiricum* Komarov, in a Prairie Wetland. *Environmental Pollution*, 95: 259-268.
25. Gardner, S.C. and C.E. Grue. 1996. Effects of Rodeo and Garlon® 3A on Non-target Wetland Species in Central Washington. *Environmental Toxicology and Chemistry* 15(4): 441-451.
26. Gardner, S.C., C.E. Grue, L.A. Grassley, J.M. Lenz, J.M. Lindenauer and M.E. Seeley. 1997. Single Species Algal (*Ankistrodesmus*) Toxicity Tests with Rodeo® and Garlon® 3A. *Bulletin of Environmental Contamination* 59: 492-499.
27. Garlon® 3A Label. Proposed in 2000.
28. Getsinger, K.D., D.G. Petty, J.D. Madsen, J.G. Skogerboe, B.A. Houtman, W.T. Haller, and A.M. Fox. 2000. Aquatic Dissipation of Herbicide Triclopyr in Lake Minnetonka, Minnesota. *Pest Manag. Sci.* 56: 388-400.
29. Getsinger, K.D., E.G. Turner, J.D. Madsen and M.D. Netherland. 1997. Restoring Native Vegetation in Eurasian Water-Milfoil Dominated Plant Community Using the Herbicide Triclopyr. *Regulated Rivers: Research & Management* 13: 357-375.
30. Getsinger, K.D. and H.E. Westerdahl. 1984. Field Evaluation of Garlon® 3A (Triclopyr and 14-ACE-F (2,4-D BEE)) for Control of Eurasian Watermilfoil. US Army Corp of Engineers, Miscellaneous Paper A-84-5.
31. Gersich, F.M., C.G. Mendoza, D.L. Hopkins, S.L. Applegath and K.M. Bodner. 1982. The Acute and Chronic Toxicity of Triclopyr ((3,5,6-Trichloro-2-pyridinyl)oxy) acetic acid, Triethylamine Salt Solution to *Daphnia magna* Straus. Dow Chemical. ES 583. MRID 0015959.
32. Gersich, F.M., C.G. Mendoza, D.L. Hopkins and K.M. Bodner. 1984. Acute and Chronic Toxicity of Triclopyr Triethylamine Salt to *Daphnia magna* Straus. *Bulletin of Environmental Contamination and Toxicology* 32:497-502.
33. Goldman, C.R and A.J. Horne, 1983. *Limnology*, McGraw-Hill Publishing Company, New York, pp. 463.

34. Gorsinske, S.J., M.A. Mayes, J.R. Ormand, J.T. Weinberg and C.H. Richardson. 1991. 3,5,6-Trichloro-2-pyridinol: Acute 96-hour Toxicity to the Bluegill, *Lepomis macrochirus* Rafinesque. Dow Chemical Company. ES-DR-0037-0423. MRID 41829003.
35. Green, W.R., H.E. Westerdahl, J.C. Joyce, and W.T. Haller. 1989. Triclopyr (Garlon® 3A) Dissipation in Lake Seminole, Georgia. Miscellaneous Paper A-89-2, US Army Engineer Waterways Experiment Station, Vicksburg, MS.
36. Hamburg, A.W., P.R. Yankovich, and B.D. Almond. 1987. The Metabolism of 14C-triclopyr in Laying Hens. ABC Laboratories and Dow Chemical. #35252. MRID 40356606.
37. Heitmuller, P.T. 1975. Acute Toxicity of M3724 to Larvae of the Eastern Oyster (*Crassostrea virginica*), Pink Shrimp (*Penaeus duroraum*) and Fiddler Crabs (*Uca pugnator*) . Bionomics – EG&G, Inc. GH-RC11; CDC235248; MRID 006263.
38. Houtman, B.A., D.R. Foster, K.D. Getsinger and D.G. Petty. 1997. Aquatic Dissipation of Triclopyr in Lake Minnetonka, Minnesota. Dow Chemical Company. ENV94001. MRID 44456102.
39. Houtman, B.A., D.R. Foster, K.D. Getsinger and D.G. Petty. 1997. Aquatic Dissipation of Triclopyr in a Whole Pond Treatment. DowElanco, ENV94012. MRID 44456103.
40. Houtman, B.A., D.R. Foster, K.D. Getsinger and D.G. Petty. 1997. Triclopyr Dissipation and Formation and Decline of its TMP and TCP Metabolites. DowElanco, ENV96052. MRID 44456107.
41. Janz, D.M., A.P. Farrell, J.D. Morgan and G.A. Vigers. 1991. Acute Physiological Stress Responses of Juvenile Coho Salmon (*Oncorhynchus Kisutch*) to Sublethal Concentrations of Garlon®4, Garlon® 3A and Vision® Herbicides. Environmental Toxicology and Chemistry. 10:73-79.
42. Lorz, H.W. and B.P. McPherson. 1976. Effects of Copper or Zinc in Fresh Water on the Adaptation to Seawater and ATPase Activity, and the Effects of Copper on Migratory Disposition of Coho Salmon (*Oncorhynchus kisutch*). J. Fish Res. Bd. Can., 33:203.
43. Lorz, H.W., S.W. Glenn, R.H. Williams, C.M. Kunkel, L.A. Norris and B.R. Loper. 1979. Effects of Selected Herbicides on Smolting Salmon. Corvallis Environmental Research Lab, OR. U.S. EPA. EPA 600/3-79-071, June, 1979.
44. Hughes, J.S. 1987. Triclopyr Triethylamine Salt: The Toxicity to *Anabaena flos-aquae*. Malcolm Pirnie, Inc. for Dow Chemical. 0460-02-1100-1. MRID 41633706.
45. Lickly, T.D. and P.G. Murphy. 1987. The Amount and Identity of [14C]Residues in Bluegills (*Lepomis macrochirus*) exposed to [14C] Triclopyr. Environmental International 13:213-218.
46. Mayes, M.A., D.C. Dill, K.M. Mendoza and K.M. Bodnar. 1983. The Acute and Chronic Toxicity of Triclopyr [(3,5,6-Trichloro-2-pyridinyl)oxy] acetic acid, Triethylamine Salt Solution to Fathead Minnows (*Pimephales promelas Rafinesque*). Dow Chemical. ES 582. MRID 0015958.
47. Mayes, M.A., D.C. Dill, K.M. Bodnar and K.M. Mendoza. 1984. Triclopyr Triethylamine Salt Toxicity to Late Stages of Fathead Minnow. Bulletin of Environmental Contamination and Toxicology. 33: 339-347.
48. McCarty, W.M. and H.C. Alexander. 1978. Toxicity of Triclopyr, Triethylamine Salt to Fresh Water Organisms. Dow Chemical. ES-199; MRID 00151956.

49. McCarty, W.M. and T.L. Batchelder. 1977. Toxicity of Triclopyr to Daphnids. Dow Chemical. ES-37L. MRID 40346504.
50. Morgan, J.D., G.A. Vigers, A.P. Farrell, D.M. Janz, and J. F. Manville. 1991. Acute Avoidance Reactions and Behavioral Responses of Juvenile Rainbow Trout (*Onchorhynchus mykiss*) to Garlon 4®, Garlon® 3A and Vision® Herbicides. Environmental Toxicology and Chemistry. 10: 73-79.
51. Neary, D.G. and J.L. Michael. 1996. Herbicides Protecting Long-Term Sustainability and Water Quality in Forest Ecosystems. New Zealand Journal of Forestry Science 26: 241-264.
52. Netherland, M.D. and K.D. Getsinger. 1993. Control of Eurasian Watermilfoil Using Triclopyr. Down to Earth 48(1): 1-5.
53. Petty, D.G., K.D. Getsinger, J.D. Madsen, J.G. Skogerboe, W.T. Haller, A.M. Fox and B.A. Houtman. 1998. Aquatic Dissipation of Herbicide Triclopyr in Lake Minnetonka, Minnesota. U.S. Army Corp of Engineers. Technical Report A-98-1.
54. Petty, D.G., J.G. Skogerboe, K.D. Getsinger, D.R. Foster, J.W. Fairchild and L.W. Anderson. 1998. Aquatic Dissipation Triclopyr in Whole-Pond Treatment. U.S. Army Corp of Engineers. Technical Report A96-6.
55. Peterson, H.G., C. Boutin, K.E. Freemark and P.A. Martin. 1994. Aquatic Phytotoxicity of 23 Pesticides Applied at Expected Environmental Concentration. Aquatic Toxicology 28:275-292.
56. Puvanesarajah, V. 1992. Metabolism of [14C]Triclopyr in Lactating Dairy Goats. ABC Laboratories, Inc. #39438. MRID 43339002.
57. Quast, J.F., et al. 1988. "Triclopyr: A One-Year Dietary Toxicity Study in Beagle Dogs". Unpublished study prepared by Mammalian and Environmental Toxicological Toxicology Research Laboratory Health and Environmental Sciences. The Dow Chemical Company. Study ID: K-042085-036.
58. Reid, G.K. 1961. Ecology of Inland Waters and Estuaries. Van Nostrand Reinhold Company, New York, NY. 375 pp.
59. Renovate EUP. 1997. Federal Experimental Use Permit, November 25, 1997.
60. Ritter, A.M. and A.L. Peacock. 2000. Aquatic Dissipation Modeling of Triclopyr. Waterborne Environmental, Inc. Dow Agro Sciences, LLC WEI 396.06. LLC 30398118-5023-1. MRID 45047901.
61. Serdar, D., A. Johnson, and M. Stinson. 1995. Seawater Challenge of Chinook Salmon Smolts (*Onchorynchus tshawytscha*) Exposed to the Aquatic Herbicide Hydrothol 191. Lab Project Number: 95-358; MRID 44700401. Unpublished study prepared by Washington State Department of Ecology. 30p.
62. Shearer, R. et al. 1996. Tacoma-Pierce County Health Department—Steilacoom Lake Phase I Restoration Study, Volume 1.
63. Shearer, R and M. Halter. 1980. Literature Reviews of Four Selected Herbicides: 2,4-D, Diclobenil, Diquat and Endothall. Metro.
64. Smith, A.E., K. Mortensen, A.J. Aubin and M.M. Molloy. 1994. Degradation of MCPA, 2,4-D and other Phenoxyalkanoic Acid Herbicides Using an Isolated Soil Bacteria. Journal of Agricultural and Food Chemistry 42(2): 401-405.

65. Swingle, H.S. 1950. Relationships and Dynamics of Balanced and unbalanced Fish Populations. Agricultural Experiment Station of the Alabama Polytechnic Institute, Bulletin No. 274.
66. Swingle, H.S. 1956. Appraisal of Methods of Fish Population Study – Part IV. Determination of Balanced Farm fish Ponds. Transactions of the 21<sup>st</sup> North American Wildlife Conference, March 5,6,7, 1956 at the Jung Hotel, New Orleans, Louisiana. Edited by J.B. Trefethen.
67. Tooby, T.E., J. Lucey and B. Scott. 1980. The Tolerance of Grass Carp, *Ctenopharyngodon idella* Val., to Aquatic Herbicides. Journal of Fish Biology 16: 591-597.
68. Urban, D.J. and N. Cook. 1986. Hazard Evaluation Division, Standard Evaluation Procedure. Ecological Risk Assessment. EPA Document Number 540/9-86/167.
69. Vedula, U., et al. 1995. “Triclopyr: Two-Generation Dietary Reproduction Study in Sprague-Dawley Rats”. Unpublished study prepared by the Toxicology Research Laboratory Health and Environmental Sciences. The Dow Chemical Company. Study ID: K-042085-048.
70. Wan, M.T., D.J. Moul and R.G. Watts. 1987. Acute Toxicity to Juvenile Pacific Salmonids of Garlon® 3A™, Garlon® 4™, triclopyr ester and their Transformation products: 3,5,6 trichloro-2-pyridinol (TCP) and 2-methoxy-3,5,6-trichloropyridine (TMP). Bulletin of Environmental Contamination and Toxicology 39: 721-728.
71. Ward, T.J. and R.L. Boeri. 1989. Acute Flow-Through Toxicity of Triclopyr TEA salt to Tidewater Silverside., *Menidia beryllina*. EnviroSystems for DowElanco. 8990-D; Dow ES-2194; MRID 41633703.
72. Ward, Y.J., P.L. Kowalski and R.L. Boeri. 1992. Acute Toxicity of Triclopyr TEA salt to the Grass Shrimp, *Palaemonetes pugio*. T.R. Wilbury for Dow Elanco. TRW 50-DO; Dow ES-2497; MRID 42646102.
73. Ward, Y.J., P.L. Kowalski and R.L. Boeri. 1993. Acute Flow-Through Mollusk Shell Deposition Test with Triclopyr TEA Salt. Dow Chemical Company. TRW 51-DO. ES-2498. MRID 42646101.
74. Woodburn, K.B. 1988. The Aquatic Dissipation of Triclopyr in Lake Seminole, Georgia. Dow Chemical. GH-C 2093. MRID 41714304.
75. Woodburn, K.B., D.D. Fontaine and J.F. Richards. 1988. A Soil Adsorption/Desorption Study with Triclopyr. Dow Chemical GH-C-2017. MRID 40749801.
76. Woodburn, K.B., W.R. Green and H.E. Westerdahl. 1993. Aquatic Dissipation of Triclopyr in Lake Seminole, Georgia. J. Agricultural and Food Chemistry 41: 2172-2177.
77. Yankovich P.R. and W.R. Bauriedel. 1976. Fate of 14C-triclopyr Fed to Lactating Goats. Dow Chemical. GH-C1017. CDC 070045F. MRID 00071805.
78. Zucker, E. 1985. Acute Toxicity Test for Freshwater Fish. Hazard Evaluation Division. Standard Evaluation Procedure. U.S. EPA. PB86-129277. EPA-540/9-85-006.

## Section 5 - Human Health Effects

1. American Conference of Governmental Industrial Hygienists (ACGHI). 1997. Threshold Limit Values (TLVs) for Chemical Substances and Physical Agents and Biological Exposure Indices (BEIs).
2. Andersen, K.J., E.G. Leighty and M.T. Takahashi. 1972. Evaluation of Herbicides for Possible Mutagenic Properties., J. Agric. Food Chem., 20:649-656.
3. Berdasco, N.M. 1990(a). 15% Dilution of Garlon 3A: Dermal Sensitization Potential in the Harley Albino Guinea Pig., Unpublished Report No. M-003724-013, Dow Chemical Company, Midland, MI, MRID 41830601.
4. Berdasco, N.M. 1990(b). 30% Dilution of Garlon 3A: Dermal Sensitization Potential in the Hartley Albino Guinea Pig., Unpublished Report No. M-003724-012, Dow Chemical Company, Midland, MI, MRID 41830602.
5. Berdasco, N.M. 1994. Four Samples of Garlon 3A: Dermal Sensitization Potential in the Hartley Albino Guinea Pig., Unpublished Report No. M-003724-011, Dow Chemical Company, Midland, MI, MRID 43230202.
6. Bryson, A.M. 1994(a). Triclopyr Triethylamine Salt: A Study of the Effect on Pregnancy of the Rat., Huntingdon Research Centre Ltd., Lab Report No. DWC 645/646/931358, DowElanco Pacific Ltd., MRID 43217602.
7. Bryson, A.M. 1994(b). Triclopyr Triethylamine Salt: A Study of the Effect on Pregnancy of the Rabbit., Huntingdon Research Centre Ltd., Lab Report No. DWC 642/641/659/931914, DowElanco Pacific Ltd., MRID 43217603.
8. Carmichael, N.G., J. Perkins, R.J. Nolan, A.P. Fletcher, R. Davies, P.M. Brown and S.J. Warrington. 1988. A Study of the Oral Absorption and Excretion of Triclopyr in Human Volunteers., Dow Chemical Co. Ltd., Agriculture Research and Development Letcomb Laboratory, Lab Project ID 87/DCS031/808, MRID 40598901.
9. Carmichael, N.G., R.J. Nolan, J.M. Perkins, R. Davies and S.J. Warrington. 1989. Oral and Dermal Pharmacokinetics of Triclopyr in Human Volunteers. Human Toxicol., 8:431-437.
10. Carmichael, N.G. 1989. Assessment of Hazards to Workers Applying Pesticides., Food Addit. Contam., 6 Suppl. 1: S21-S27.
11. Chakrabarti, A. and S. Gennrich. 1988. Attempted Determination of the Vapor Pressure of Triclopyr TEA Salt by the Knudsen-Effusion/Weight Loss Method: Project ID: ML-AL 88-020173. Unpublished Report. Dow Chemical Company.
12. Dow AgroSciences. 2000. Personal Communication with Dr. J.R. Peterson, Manager of Product Registration.
13. Dow Chemical Company, North American Agricultural Products Department. 1988. Environmental Fate Summary Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid).

14. Eisenbrandt, D.L., H.M. Firchau, E.L. Wolfe and T.D. Landry. 1987. Triclopyr: 2-Year Dietary Chronic Toxicity-Oncogenicity Study in Fischer 344 Rats. HET K-04085-026. Health and Environmental Sciences. Dow Chemical Company, Midland, MI. MRID 40107701.
15. Eisenbrandt, D.L., R.J. Nolan, S.A. McMaster and L.H. McKendry. 1997. Triclopyr: An Assessment of Common Mechanism of Toxicity. Dow Chemical Company. Midland, MI. Unpublished Report No. HET K-o42085-097. MRID 44385901.
16. Fabrizio, D.P.A. 1973(a). Dominant Lethal Assay for Mutagenesis: Compound Dowco 233: LBI Project No. 2421. Final Report Litton Bionetics, Inc. for Dow Chemical Company, Midland, MI. MRID 00057087.
17. Fabrizio, D.P.A. 1973b. Acute and Subacute in vivo Host-Mediated Assay for Mutagenesis: Compound Dowco 233: LBI Project No. 2421. Final Report Litton Bionetics, Inc. for Dow Chemical Company, Midland, MI. MRID 00057086.
18. Feldman, R.J. and H.I. Maibach. 1974. Percutaneous Penetration of Some Pesticides and Herbicides in Man. *Toxicol. Appl. Pharmacol.*, 29:126-132.
19. Frank, P.A. and R.D. Comes. 1967. Herbicidal Residues in Pond Water and Hydrosol. *Weeds*, 15:210-213.
20. GARLON® 3A Label. 2000. DowAgroSciences LLC.
21. Getsinger, K.D., et al. 1977. Restoring Native Vegetation in a Eurasian Watermilfoil Dominated Plant Community Using the Herbicide Triclopyr. *Regulated Rivers: Res. and Management*. 13:357-375.
22. Gilbert, K.S. 1996. Garlon® 3A: Acute Dermal Toxicity Study in New Zealand White Rabbits. Unpublished Report No. M-003724-015D, Dow Chemical Company, Midland, MI. MRID 43952401.
23. Goodman, D.G. and P.K. Hildebrandt. 1988. Review of Tumor Data From the Two-Year Dietary Chronic Toxicity/Oncogenicity Study on Triclopyr in F344 Rats. Lab Project No. MLJ61495, PATHCO, Inc., Ijamsville, MD for Dow Elanco. MRID 43683401.
24. Goodman, D.G. and P.K. Hildebrandt. 1996. Review of Tumor Data From the Chronic Toxicity/Carcinogenicity Studies of Triclopyr in F344 Rats and ICR(Jcl:ICR) Mice. Lab Project No. PATHCO No. 96-86. PATHCO, Inc., Ijamsville, MD for DowElanco,. MRID 44302501.
25. Grzenda, A.R., H.P. Nicholson and W.S. Cox. 1966. Persistence of Four Herbicides in Pond Water. *J. Am. Water Works Ass.*, 58:326-332.
26. Hanley, T.R., D.J. Thompson, A.K. Palmer, R.P. Beliles and B.A. Schwetz. 1984. Teratology and Reproduction Studies with Triclopyr in the Rat and Rabbit. *Fundam. Appl. Toxicol.*, 4: 872-882.
27. Henck, J.W. 1980(a). Garlon® 3A Herbicide/Water Solution (1:3) Eye and Skin Irritation Potential. Dow Chemical Company, Midland, MI. MRID 00086799.

28. Henck, J.W. 1980(b). Garlon 3A Herbicide/Water Solution (1:7) Eye and Skin Irritation Potential., Dow Chemical Company, Midland, MI, MRID 00086800.
29. Hotchkiss, S.A., P. Hewitt, J. Caldwell, W.L. Chen and R.R Rowe. 1992 Percutaneous Absorption of Nicotinic Acid, Phenol, Benzoic Acid and Triclopyr Butoxyethyl Ester Through Rat and Human Skin *in vitro*: Further Validation of an *in vitro* model by comparison with *in vivo* Data. Food Chem. Toxicol., 30:891-899.
30. Landry, T.D., D. Eisenbrandt and T. Gushow. 1984(a). Triclopyr: 13-Week Dietary Toxicity in Fischer 344 Rats. Unpublished Report, Dow Chemical Company, Midland, MI. MRID 00150378.
31. Landry, T.D., P.E. Kastl and T.S. Gushow. 1984(b). Triclopyr: Dietary Pharmacokinetics Study in Fischer 344 Rats. Dow Chemical Company, Midland, MI. MRID 00150379.
32. Levin, D.E., M. Hollstein, M.F. Christman, E.A. Schwiers and B.N. Ames. 1982. A New Salmonella Tester Strain (TA 102) with AT Base Pairs at the Site of Mutation Detects Oxidative Mutagens. Proc. Nat. Acad. Sci., 79:7445-7449.
33. Lockwood, D.D. 1992. Garlon® 3A Herbicide: Dermal Probe Study in New Zealand White Rabbits. Unpublished Report No. M-003724-014, Dow Chemical Company, Midland, MI. MRID 42339001.
34. Lunchick, C. 1994. Assessment of Swimmer Exposure to Endothall in Aquathol®-and Hydrothol®-Treated Bodies of Water. Report Prepared by: Jellinek, Schwartz & Connolly, Inc. Arlington, VA. Report to Elf Atochem North America, Inc
35. McMahon, C.K. and P.B. Bush. 1992. Forest Worker Exposure to Airborne Herbicide Residues in Smoke From Prescribed Fires in the Southern United States. Am. Ind. Hyg. Assoc. J., 53:265-272.
36. Mendrala, A. and M. Dryzga. 1986. The Evaluation of Triclopyr in the Rat Hepatocyte Unscheduled DNA Synthesis Assay: Laboratory Project ID; HET K-042085-031. Dow Chemical Company, Midland, MI. MRID 40055702.
37. Merck Index; An Encyclopedia of Chemicals, Drugs, and Biologicals. 1989. Editor: Susan Budavari, Eleventh Edition, Published by Merck & Co., Inc., P. 9572-9573.
38. Middendorf, P., C. Timchalk, B. Kropscott and D. Rick. 1994. Forest Worker Exposure to Garlon® 4 Herbicide., Appl. Occup. Environ. Hyg., 9:589-594.
39. Mizell, M.J. and L.G. Lomax. 1988(a). Garlon® 3A (Triclopyr as Triethylamine Salt): Acute Oral Toxicity Study in Fischer 344 Rats. Unpublished Report No. M-003724-009A, Dow Chemical Company, Midland, MI. MRID 41443301.
40. Mizell, M.J. 1988(b). Garlon® 3A (Triclopyr as Triethylamine Salt): Primary Dermal Irritation Study in New Zealand White Rabbits. Unpublished Report No. M-003724-009B, Dow Chemical Company, Midland, MI. MRID 41443305.



41. Mizell, M.J. 1988(c). Garlon® 3A (Triclopyr as Triethylamine Salt): Primary Eye Irritation Study in New Zealand White Rabbits. Unpublished Report No. M-003724-009C, Dow Chemical Company, Midland, MI. MRID 41443304.
42. Mizell, M.J. 1989. Garlon® 3A (Triclopyr as Triethylamine Salt): Acute Dermal Toxicity Study in New Zealand White Rabbits. Unpublished Report No. M-003724-009D, Dow Chemical Company, Midland, MI. MRID 41443302.
43. Mizell, M.J. 1989. Garlon® 3A (Triclopyr as Triethylamine Salt): Dermal Sensitization Potential in the Hartley Albino Guinea Pig. Unpublished Report No. M-003724-009E, Dow Chemical Company, Midland, MI. MRID 41443306.
44. Montgomery, J.H. 1993. Agrochemicals Desk Reference, Environmental Data. Lewis Publishers, Boca Raton, FL, pp. 180-181.
45. Nitschke, K.D., J.E. Battjes and B.L. Yano. 1989. XRM-3724: Acute Aerosol LC50 Study in Fischer 344 Rats. Unpublished Report No. M-003724-010, Dow Chemical Company, Midland, MI. MRID 41443303.
46. Petty, D.G. et al. 1998. Aquatic Dissipation of the Herbicide Triclopyr in Lake Minnetonka, Minnesota. Technical Report A-98-1. U.S. Army Corps of Engineers.
47. Quast, J.F., C.G. Humiston and J.E. LeBeau. 1976. 3,5,6- Trichloro-2-pyridyloxyacetic Acid (Dowco 233 Herbicide): Subchronic Dietary Feeding Study in Beagle Dogs. Unpublished Study Report No. CDL:0700043-C, Dow Chemical Company, Midland, MI., MRID 00071793.
48. Quast, J.F., C.E. Wade and R.V. Kalnins. 1977. 3,5,6-Trichloro-2-pyridinyloxyacetic Acid (Dowco 233 Herbicide): Supplemental Subchronic Dietary Feeding Study in Beagle Dogs. Unpublished Study Report No. CDL:070043-D., Dow Chemical Company, Midland, MI. MRID 00071794.
49. Quast, J.F., T.S. Gushow, W.T. Stott, J.M. Wall and B.D. Landenberger. 1988. Triclopyr: A One-Year Dietary Toxicity Study in Beagle Dogs. Unpublished Study Report No. K-042085-036, Dow Chemical Company, Midland, MI. MRID 41200301.
50. Rao, K.S., T.R. Hanley, Jr. and B. Dabny. 1979. Dowco 233: Heritable Translocation and Dominant Lethal Test in the Mouse. Unpublished Report CDL:241822-A, Dow Chemical Company, Midland, MI. MRID 00028996.
51. Renovate® Label. 1997. DowElanco.
52. Richold, M., E. Jones and R.J. Proudlock. 1979. Ames Metabolic Activation Test to Assess the Potential Mutagenic Effect of Dowco 233: DWC316/791014. Unpublished Report CDL:242367-D, Dow Chemical Company, Midland, MI. MRID 00031939.
53. Shipp, A.M. 1986. Worst Case Analysis Study on Forest Plantation Herbicide Use. Prepared for the Forest Land Management Division, Department of Natural Resources, State of Washington.

54. Shirasu, Y., M. Moriya and K. Watanabe. 1978. Report of Mutagenicity Test on Triclopyr in Bacterial Test System. Prepared by Institute of Environmental Toxicology, Japan, Report No. CDL:242367-C, Dow Chemical Company, Midland, MI. MRID 00038408.
55. Sibinovic, K.H. 1973. *In Vitro* and Subacute *In Vivo* Host-Mediated Assay for Mutagenesis: Compound Dowco 233: LBI Project No. 2421., Prepared by Litton Bionetics, Inc. Dow Chemical Company, Midland, MI. MRID 00057085.
56. Smith, F.A., K.A. Brzak, G.A. Bormett and R.J. Nolan. 1992. Triclopyr, Triethylamine Salt: Pharmacokinetics and Metabolism in Male Fischer 344 Rats. Dow Chemical Company Midland, MI. Unpublished Report DR-0114-2861-003. MRID 42437901 and 42444701.
57. Solomon, K.R. et al. 1988. Persistence of Hexazinon (Velpar), Triclopyr (Garlon), and 2,4-D in a Northern Ontario Aquatic Environment. *J. Agric. Food Chem.*, 36:1314-1318.
58. Timchalk, C., M.D. Dryzga and P.E. Kastl. 1988. Triclopyr: Tissue Distribution and Metabolism of <sup>14</sup>C-Labeled Triclopyr in Fischer 344 Rats. Dow Chemical Company, Midland, MI. Report No. K-42085-(40). MRID 41353001.
59. Timchalk, C., M.D. Dryzga and P.E. Kastl. 1990. Pharmacokinetics and Metabolism of Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid) in Fischer 344 Rats. *Toxicology*, 62:71-87.
60. Timchalk, C., D.R. Finco and J.F. Quast. 1997(a). Evaluation of Renal Function in Rhesus Monkeys and Comparison to Beagle Dogs Following Oral Administration of the Organic Acid Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid). *Fund. Appl. Toxicol.* 36:47-53.
61. Timchalk, C. and R.J. Nolan. 1997(b). Pharmacokinetics of Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid) in the Beagle Dog and Rhesus Monkey: Perspective on the Reduced Capacity of Dogs to Excrete the Organic Acid Relative to the Rat, Monkey and Human. *Toxicol. Appl. Pharmacol.*, 144:268-278.
62. Tsuda, S., K. Ebino, M. Ikeda, T. Harada and Y. Shirasu. 1987. Triclopyr: 22-Month Oral Chronic Toxicity and Oncogenicity Study in Mice. GHF-R-111. The Institute of Environmental Toxicology, Mitseikaido, Japan for Dow Chemical, Japan, Ltd. MRID 40356601.
63. U.S. EPA. 1987. Reference Dose (RfD): Description and Use in Health Risk Assessments. Integrated Risk Information System, Appendix A. Intra-Agency Reference Dose Work Group, Environmental Criteria and Assessment Office, Cincinnati, OH.
64. US EPA 1989. Risk Assessment Guidance for Superfund. Vol. I Human Health Evaluation Manual (Part A). EPA/540/1.
65. US EPA. 1998. Reregistration Eligibility Decision (RED) Triclopyr. Prevention, Pesticides and Toxic Substances (7508C), EPA 738-R-98-011.
66. van Beck, L. and D.C. Leegwater. 1981(a). Study on the Percutaneous Absorption of Triclopyr by the Rabbit. Part II: Single Treatment of Intact Skin. Civo Institutes TNO, Report No. GHE-T-033, Zeist, Netherlands, Dow Chemical Company. MIRD 00153805.

67. van Beck, L., D.C. Leegwater and J.B. Bruyntjes. 1981(b). Subacute (3-Week) Percutaneous Absorption Study with Garlon® 4E in Albino Rabbits., Part I: Repeated Treatment with 0.5 and 1.0 ml 50% Aqueous Garlon® 4E/kg Body Weight. Civo Institutes TNO, Report No. GHE-T-043, Zeist, Netherlands, Dow Chemical Company, MIRD 00153846.
68. van Beck, L., H.E. Falke and M.C. Bosland. 1984. Subacute (3-Week) Dermal Toxicity Study with Garlon 4 in Rats (Final Report). CIVO Institutes TNO, Zeist, Netherlands, Unpublished Report No. DET 408. Submitted by Dow Chemical Company 9/30/85 under EPA No. 464-554. MRID 0015346.
69. Vedula, U., W.J. Breslin, B.E. Kropscott and K.E. Stebbins. 1995. Triclopyr: Two-Generation Dietary Reproduction Study in Sprague-Dawley Rats. Study Report No. K-042085-048, DowElanco. MRID 43545701.
70. Washington Department of Health. 1999. Review of Proposed Spot Treatment with Renovate Aquatic Herbicide.
71. Wester, R.C. and H.I. Maibach. 1985. In Vivo Percutaneous Absorption and Decontamination of Pesticides in Humans. *J. Toxicol. Environ. Health*, 16:25-37.
72. Woodburn, K.B. et al. 1993(a). Aquatic Photolysis of Triclopyr., *Environ. Toxicol. Chem.*, 12:43-55.
73. Woodburn, K.B. et al. 1993(b). Aquatic Dissipation of Triclopyr in Lake Seminole, Georgia. *J. Agric. Food Chem.*, 41:2172-2177.
74. Worthing, C.R. and R.J. Hance. 1991. *The Pesticide Manual*, 9th Edition.